

Homework 2 - Einstein equation

Q2.1. Use the Bianchi identity

$$\nabla_{\mathbf{a}} R_{\mathbf{bcd}}^{\mathbf{e}} + \nabla_{\mathbf{b}} R_{\mathbf{cad}}^{\mathbf{e}} + \nabla_{\mathbf{c}} R_{\mathbf{abd}}^{\mathbf{e}} = 0 \quad (\text{Q2.1.1})$$

to show that

$$\nabla_{\mathbf{b}} G_{\mathbf{a}}^{\mathbf{b}} = 0 \quad (\text{Q2.1.2})$$

A2.1. Contracting the \mathbf{c} and \mathbf{e} indices, contracting the \mathbf{a} and \mathbf{d} indices using the metric, using the symmetries Eqs. (1.2.8) and (1.2.9) and the definitions Eqs (1.2.12), (1.2.13) and (2.4.7) gives Eq. (Q2.1.2)

$$0 = g^{\mathbf{ad}} \nabla_{\mathbf{a}} R_{\mathbf{bcd}}^{\mathbf{c}} + g^{\mathbf{ad}} \nabla_{\mathbf{b}} R_{\mathbf{cad}}^{\mathbf{c}} + g^{\mathbf{ad}} \nabla_{\mathbf{c}} R_{\mathbf{abd}}^{\mathbf{c}} \quad (\text{A2.1.1})$$

$$= \nabla_{\mathbf{a}} R_{\mathbf{bc}}^{\mathbf{ac}} - \nabla_{\mathbf{b}} R_{\mathbf{ac}}^{\mathbf{ac}} + \nabla_{\mathbf{c}} R_{\mathbf{ba}}^{\mathbf{ca}} \quad (\text{A2.1.2})$$

$$= 2\nabla_{\mathbf{a}} R_{\mathbf{b}}^{\mathbf{a}} - \nabla_{\mathbf{b}} R \quad (\text{A2.1.3})$$

$$= 2\nabla_{\mathbf{a}} G_{\mathbf{b}}^{\mathbf{a}} \quad (\text{A2.1.4})$$

Q2.2. Calculate

$$\nabla_{\mathbf{b}} T_{\mathbf{a}}^{\mathbf{b}} = 0 \quad (\text{Q2.2.1})$$

for a perfect fluid and interpret your answer.

A2.2. Eq. (2.4.13) gives

$$0 = \nabla_{\mathbf{b}} T_{\mathbf{a}}^{\mathbf{b}} = \nabla_{\mathbf{b}} [\rho u_{\mathbf{a}} u^{\mathbf{b}} + p (u_{\mathbf{a}} u^{\mathbf{b}} - \delta_{\mathbf{a}}^{\mathbf{b}})] \quad (\text{A2.2.1})$$

$$= u_{\mathbf{a}} u^{\mathbf{b}} \nabla_{\mathbf{b}} \rho + (\rho + p) u^{\mathbf{b}} \nabla_{\mathbf{b}} u_{\mathbf{a}} + (\rho + p) u_{\mathbf{a}} \nabla_{\mathbf{b}} u^{\mathbf{b}} + (u_{\mathbf{a}} u^{\mathbf{b}} - \delta_{\mathbf{a}}^{\mathbf{b}}) \nabla_{\mathbf{b}} p \quad (\text{A2.2.2})$$

therefore, using Eqs. (2.3.3) and (2.3.4),

$$u^{\mathbf{a}} \nabla_{\mathbf{b}} T_{\mathbf{a}}^{\mathbf{b}} = u^{\mathbf{b}} \nabla_{\mathbf{b}} \rho + (\rho + p) \nabla_{\mathbf{b}} u^{\mathbf{b}} = 0 \quad (\text{A2.2.3})$$

and

$$(g^{\mathbf{ac}} - u^{\mathbf{a}} u^{\mathbf{c}}) \nabla_{\mathbf{b}} T_{\mathbf{c}}^{\mathbf{b}} = (\rho + p) u^{\mathbf{b}} \nabla_{\mathbf{b}} u^{\mathbf{a}} + (u^{\mathbf{a}} u^{\mathbf{b}} - g^{\mathbf{ab}}) \nabla_{\mathbf{b}} p = 0 \quad (\text{A2.2.4})$$

which are the relativistic Euler equations, corresponding to energy and momentum conservation in the rest frame of the fluid. Writing the energy density ρ in terms of the mass density ρ_m and non-mass energy per unit mass e

$$\rho = \rho_m (1 + e) \quad (\text{A2.2.5})$$

Eq. (A2.2.3) gives

$$(1 + e) (u^{\mathbf{b}} \nabla_{\mathbf{b}} \rho_m + \rho_m \nabla_{\mathbf{b}} u^{\mathbf{b}}) + (\rho_m u^{\mathbf{b}} \nabla_{\mathbf{b}} e + p \nabla_{\mathbf{b}} u^{\mathbf{b}}) = 0 \quad (\text{A2.2.6})$$

corresponding to conservation of mass and Newtonian energy.